PubMed

Format: Abstract

Full text links

ELSEVIER FULL-TEXT ARTICLE

Exp Mol Pathol. 2002 Aug;73(1):19-27.

Antibodies to squalene in recipients of anthrax vaccine.

Asa PB[1], Wilson RB, Garry RF.

Author information

Abstract

We previously reported that antibodies to squalene, an experimental vaccine adjuvant, are present in persons with symptoms consistent with Gulf War Syndrome (GWS) (P. B. Asa et al., Exp. Mol. Pathol 68, 196-197, 2000). The United States Department of Defense initiated the Anthrax Vaccine Immunization Program (AVIP) in 1997 to immunize 2.4 million military personnel. Because adverse reactions in vaccinated personnel were similar to symptoms of GWS, we tested AVIP participants for anti-squalene antibodies (ASA). In a pilot study, 6 of 6 vaccine recipients with GWS-like symptoms were positive for ASA. In a larger blinded study, only 32% (8/25) of AVIP personnel compared to 15.7% (3/19) of controls were positive ($P > 0.05$). Further analysis revealed that ASA were associated with specific lots of vaccine. The incidence of ASA in personnel in the blinded study receiving these lots was 47% (8/17) compared to an incidence of 0% (0/8; $P < 0.025$) of the AVIP participants receiving other lots of vaccine. Analysis of additional personnel revealed that in all but one case (19/20; 95%), ASA were restricted to personnel immunized with lots of vaccine known to contain squalene. Except for one symptomatic individual, positive clinical findings in 17 ASA-negative personnel were restricted to 4 individuals receiving vaccine from lots containing squalene. ASA were not present prior to vaccination in preimmunization sera available from 4 AVIP personnel. Three of these individuals became ASA positive after vaccination. These results suggest that the production of ASA in GWS patients is linked to the presence of squalene in certain lots of anthrax vaccine.

PMID: 12127050

[Indexed for MEDLINE]

Publication types, MeSH terms, Substances

LinkOut - more resources